

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 967 205 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
29.12.1999 Bulletin 1999/52

(51) Int Cl.<sup>6</sup>: **C07D 213/38**, **C07D 401/14**,  
**G01N 33/48**

(21) Application number: **99660100.1**

(22) Date of filing: **03.06.1999**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: **25.06.1998 US 104219**

(71) Applicant: **WALLAC OY**  
**20101 Turku (FI)**


(72) Inventors:  
• **Takalo, Harri**  
**20360 Turku (FI)**

- **Hovinen, Jari**  
**20320 Turku (FI)**
- **Mukkala, Veli-Matti**  
**20780 Kaarina (FI)**
- **Liitti, Päivi**  
**20750 Turku (FI)**
- **Mikola, Heikki**  
**20720 Turku (FI)**

(74) Representative: **Öhman, Ann-Marie**  
**Turun Patenttitoimisto Oy,**  
**P.O. Box 99**  
**20521 Turku (FI)**

**(54) Labeling reactants and their use**

(57) The invention relates to a novel labeling reactant, suitable for labeling of a biospecific binding reactant using solid-phase synthesis. The invention further concerns new labeling methods. The novel labeling reactant has the formula (I)

**(I)**

wherein

**A** is a bivalent aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate;

**R** is - Z(G' - NH - X) G" - E, where

**X** is a transient protecting group;

**E** is a carboxylic acid, its salt, active ester or halide;

**Z** is the bridge point;

**G** is a bridge between A and Z, **G'** is a bridge between NH and Z and **G"** is a bridge between E and Z.

**R'** is -COOR''' where R''' is an alkyl of 1 to 4 carbon atoms, phenyl or benzyl.

**EP 0 967 205 A1**

## Description

## FIELD OF THE INVENTION

**[0001]** This invention relates to novel compounds suitable for labeling of a biospecific binding reactant using solid-phase synthesis. The invention relates further to novel methods of labeling a biospecific binding reactant in a solid-phase synthesis.

## BACKGROUND OF THE INVENTION

**[0002]** The publications and other materials used herein to illuminate the background of the invention, and in particular, cases to provide additional details respecting the practice, are incorporated by reference.

**[0003]** Solid-phase synthesis is a well-known and widely spread technology in modern organic chemistry for preparation of various biologically important compounds. It has found many new applications such as combinatorial libraries and the peptide-based DNA analogue (PNA, peptide nucleic acids). Along with the technology development, the need for labeling those biologically interesting compounds during solid-phase synthesis has appeared. Several non-fluorescent labeling candidates have been published (Arya, R. and Gariépy, J., 1991, *Bioconjugate Chem.*, 2, 323, Cuenoud, B. and Schepartz, A., 1991, *Tetrahedron*, 47, 2535; Rana, T.M., Ban, M. and Hearst, J.E., 1992, *Tetrahedron Lett.*, 33, 4521; Song, A.I. and Rana, T.A., 1997, *Bioconjugate Chem.*, 8, 249). These labels have been developed mainly to incorporate metal ions such as Fe, Cu in a peptide to act as sequence-specific cleaving agents for proteins. Radioactive  $^{153}\text{Gd}$ ,  $^{111}\text{In}$  etc. can be used for in vivo tumor localization and radioimmunotherapy. These published monomers can be used as luminescent labels using lanthanide ions and a separate luminescence enhancement step but a separate enhancement step is needed to produce the luminescence to be detected. During the luminescence enhancement the lanthanide has to be dissociated from the label and then spatial information will be lost.

**[0004]** Fluorescent label monomers for solid-phase synthesis have been published (Lohse, J., Nielson, P.E., Harrit, N. and Dahl, O., 1997, *Bioconjugate Chem.*, 8, 503; McCafferty, D.G., Bishop, B.M., Wall, C.G., Hughes, S.G., Mecklenberg, S.L., Meyer, T.J. and Erickson, B.W., 1995, *Tetrahedron*, 51, 1093; WO 96/03409). Fluorescein and other organic chromophores were used in those studies. However, such labels and labeled biomolecules suffer from many commonly known drawbacks such as Raman scattering, other fluorescent impurities, low water solubility, concentration quenching etc.

**[0005]** In the specific binding assays, such as e.g. immunoassays, DNA hybridization assays, receptor-binding assays, and cellular binding assays, generally very low concentrations of the analytes to be measured are present. Therefore, various labeling compounds have been developed that allow the labeled reactant to be detected and quantitated with high sensitivity. In immunoassays and DNA hybridization assays, time-resolved luminescence spectroscopy using lanthanide chelates is well known (e.g. I. Hemmilä, T. Ståhlberg, and P. Moltram (eds.), "Bioanalytical Applications of Labeling Technologies", Wallac, Turku, 1994). Stable photoluminescent (in this context simply referred to as luminescent) lanthanide chelates also have other applications, e.g. fluorescence microscopy and cytometry. Therefore, a number of attempts have been made to develop new highly luminescent chelate labels suitable for those types of time-resolved fluorometric applications. These include e.g. stable chelates composed of derivatives of pyridines (US 4,920,195, US 4,801,722, US 4,761,481, PCT/FI91/00373, US 5,459,186, EP Appl. 0770610, Remuñán, M.J., Roman, H., Alonso, M.T. and Rodríguez-Ubis, J.C., 1993, *J. Chem. Soc., Perkin Trans.2*, 1099), bipyridines (US 5,216,134), terpyridines (US 4,859,777, US 5,202,423, US 5,324,825) or various phenolic compounds (US 4,670,572, US 4,794,191, Ital. Pat. 42508 A/89) as the energy mediating groups and polycarboxylic acids as chelating parts. In addition, various dicarboxylate derivatives (US 5,032,677, US 5,055,578, US 4,772,563), macrocyclic cryptates (US 4,927,923, WO 93/5049, EP-A-493 745) and macrocyclic Schiff bases (EP-A-369 000) have been patented. None of these patents and articles present any methods for labeling of a biospecific binding reactant such as a hapten, a peptide, a receptor ligand, a drug or PNA oligomer, with luminescent labels by using solid-phase synthesis.

## OBJECTS AND SUMMARY OF THE INVENTION

**[0006]** A main purpose of the present invention is to solve the problems associated with labeling of small molecules with luminescent chelates and to develop easier labeling methods. By preparing suitable ligand structures together with solid-phase synthesis the labeling of small molecules with directly luminescent lanthanide chelates is achievable.


**[0007]** One object of this invention is to provide a novel labeling reactant suitable for labeling of a biospecific binding reactant using solid-phase synthesis.

**[0008]** Another object of the present invention is to provide a novel method for labeling of a biospecific binding reactant in solid-phase synthesis

**[0009]** Yet another object of the present invention is to provide a method of labeling a biospecific binding reactant

with an exact, predetermined number of labeling reactants (one or several)

[0010] Thus according to one aspect, the present invention concerns a novel labeling reactant of formula (I) suitable for labeling of a biospecific binding reactant using solid-phase synthesis



wherein


A is a bivalent aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate;

R is -Z(G'-NH-X)G"-E, where

X is a transient protecting group, where said group is e.g. 2-(4-nitro-phenylsulfonyl)ethoxycarbonyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, *tert*-butoxycarbonyl, fluorenylmethyloxycarbonyl;

E is a carboxylic acid, its salt, active ester or halide, where the said ester is e.g. an *N*-hydroxysuccinimido, *o*-, *p*- or *m*-nitrophenol, 2,4-dinitrophenol or pentafluorophenol, and the said halide is fluoride, bromide or chloride;

Z is the bridge point and is formed from




or trivalent derivatives, substituted or unsubstituted, of cyclohexane, cyclohexene, cyclohexadiene, benzene, cyclopentane, cyclopentene, cyclopentadiene, cyclobutane, cyclobutene, cyclobutadiene, aziridine, diaziridine, oxetane, thietane, azete, azetidene, 1,2-dihydro-1,2-diazete, 1,2-diazetidene, furan, tetrahydrofuran, thiophene, 2,5-dihydrothiophene, thiolane, selenophene, pyrrole, pyrrolidine, phosphole, 1,3-dioxolane, 1,2-dithiole, 1,2-thiolane, 1,3-dithiole, 1,3-dithiolane, oxazole, 4,5-dihydrooxazole, isoxazole, 4,5-dihydroisoxazole, 2,3-dihydroisoxazole, thiazole, isothiazole, imidazole, imidazolidine, pyrazole, 4,5-dihydropyrazole, pyrazolidine, triazole, pyran, pyran-2-one, 3,4-dihydro-2H-pyran, tetrahydropyran, 4H-pyran, pyran-4-one, pyridine, pyridone, piperidine, phosphabenzene, 1,4-dioxin, 1,4-dithiin, 1,4-oxathiin, oxazine, 1,3-oxazinone, morpholine, 1,3-dioxane, 1,3-dithiane, pyridazine, pyrimidine, pyrazine, piperazine, 1,2,4-triazine, 1,3,5-triazine, 1,3,5-triaza-cyclohexane-2,4,6-trione;

G is a bridge between A and Z. G' is a bridge between NH and Z and G" is a bridge between E and Z, same or different, not necessary present, and are formed of one to ten moieties, each moiety being selected from the group consisting of phenylene, alkylene containing 1-12 carbon atoms, ethynediyl (-C≡C-), ether (-O-), thioether (-S-), amide (-CO-NH- and -NH-CO- and -CO-NR'- and -NR'-CO-), carbonyl (-CO-), ester (-COO- and -OOC-), disulfide (-S-S-), diaza (-N=N-), and tertiary amine (-N(R')-), wherein R' represents an alkyl containing less than 5 carbon atoms;

R' is -COOR''' where R''' is an alkyl of 1 to 4 carbon atoms, phenyl or benzyl, which phenyl or benzyl can be substituted or unsubstituted

[0011] According to another aspect, this invention concerns a novel method of labeling a biospecific binding reactant in a solid-phase synthesis. Said method comprises the steps of

a) reacting a labeling reactant of formula (I)



wherein


A is a bivalent aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate;

R is -Z(G'-NH-X)G"-E, where

X is a transient protecting group, where said group is e.g. 2-(4-nitrophenylsulfonyl)ethoxycarbonyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, *tert*-butoxycarbonyl, fluorenylmethyloxycarbonyl;

E is a carboxylic acid, its salt, active ester or halide, where the said ester is e.g. an *N*-hydroxysuccinimido, *o*-, *p*- or *m*-nitrophenol, 2,4-dinitrophenol or pentafluorophenol, and the said halide is fluoride, bromide or chloride;

Z is the bridge point and is formed from



or trivalent derivatives, substituted or unsubstituted, of cyclohexane, cyclohexene, cyclohexadiene, benzene, cyclopentane, cyclopentene, cyclopentadiene, cyclobutane, cyclobutene, cyclobutadiene, aziridine, diaziridine, oxetane, thietane, azete, azetidine, 1,2-dihydro-1,2-diazete, 1,2-diazetidene, furan, tetrahydrofuran, thiophene, 2,5-dihydrothiophene, thiolane, selenophene, pyrrole, pyrrolidine, phosphole, 1,3-dioxolane, 1,2-dithiole, 1,2-thiolane, 1,3-dithiole, 1,3-dithiolane, oxazole, 4,5-dihydrooxazole, isoxazole, 4,5-dihydroisoxazole, 2,3-dihydroisoxazole, thiazole, isothiazole, imidazole, imidazolidine, pyrazole, 4,5-dihydropyrazole, pyrazolidine, triazole, pyran, pyran-2-one, 3,4-dihydro-2H-pyran, tetrahydropyran, 4H-pyran, pyran-4-one, pyridine, pyridone, piperidine, phosphabenzene, 1,4-dioxin, 1,4-dithiin, 1,4-oxathiin, oxazine, 1,3-oxazinone, morpholine, 1,3-dioxane, 1,3-dithiane, pyridazine, pyrimidine, pyrazine, piperazine, 1,2,4-triazine, 1,3,5-triazine, 1,3,5-triaza-cyclohexane-2,4,6-trione;

G is a bridge between A and Z, G' is a bridge between NH and Z and G" is a bridge between E and Z, same or different, not necessary present, and are formed of one to ten moieties, each moiety being selected from

the group consisting of phenylene, alkylene containing 1-12 carbon atoms, ethynediyl ( $-C\equiv C-$ ), ether ( $-O-$ ), thioether ( $-S-$ ), amide ( $-CO-NH-$  and  $-NH-CO-$  and  $-CO-NR'-$  and  $-NR'-CO-$ ), carbonyl ( $-CO-$ ), ester ( $-COO-$  and  $-OOC-$ ), disulfide ( $-S-S-$ ), diaza ( $-N=N-$ ), and tertiary amine ( $-N(R'')-$ ), wherein  $R''$  represents an alkyl containing less than 5 carbon atoms:

$R'$  is  $-COOR'''$  where  $R'''$  is an alkyl of 1 to 4 carbon atoms, phenyl or benzyl, which phenyl or benzyl can be substituted or unsubstituted,

with a functional group which is bound to a solid phase,


b) cleaving the product obtained in step a) from the solid phase, converting  $R'$  to  $-COOH$ , and removing optional protection groups,

c) converting the product from step b) to a lanthanide chelate, and

d) in case the functional group mentioned in step a) is not part of the final biospecific binding specific reactant, attaching said biospecific binding reactant to the lanthanide chelate obtained in step c).

**[0012]** According to yet another aspect, this invention concerns a novel method of labeling a biospecific binding reactant with an exact, predetermined number of labeling reactants of formula (I) in a solid-phase synthesis, said method comprising the steps of

a) reacting a labeling reactant of formula (I)



wherein


**A** is a bivalent aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate;

**R** is  $-Z(G'-NH-X)G''-E$ , where

**X** is a transient protecting group, where said group is e.g. 2-(4-nitrophenylsulfonyl)ethoxycarbonyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, *tert*-butoxycarbonyl, fluorenylmethyloxycarbonyl;

**E** is a carboxylic acid, its salt, active ester or halide, where the said ester is e.g. an *N*-hydroxysuccinimido, *o*-, *p*- or *m*-nitrophenol, 2,4-dinitrophenol or pentafluorophenol, and the said halide is fluoride, bromide or chloride;

**Z** is the bridge point and is formed from



or trivalent derivatives, substituted or unsubstituted, of cyclohexane, cyclohexene, cyclohexadiene, benzene, cyclopentane, cyclopentene, cyclopentadiene, cyclobutane, cyclobutene, cyclobutadiene, aziridine, diaziridine, oxetane, thietane, azete, azetidine, 1,2-dihydro-1,2-diazete, 1,2-diazetidine, furan, tetrahydrofuran, thiophene, 2,5-dihydrothiophene, thiolane, selenophene, pyrrole, pyrrolidine, phosphole, 1,3-dioxolane, 1,2-dithiole, 1,2-thiolane, 1,3-dithiole, 1,3-dithiolane, oxazole, 4,5-dihydrooxazole, isoxazole, 4,5-dihydroisoxazole, 2,3-dihydroisoxazole, thiazole, isothiazole, imidazole, imidazolidine, pyrazole, 4,5-dihydropyrazole, pyrazolidine, triazole, pyran, pyran-2-one, 3,4-dihydro-2H-pyran, tetrahydropyran, 4H-pyran, pyran-4-one, pyridine, pyridone, piperidine, phosphabenzene, 1,4-dioxin, 1,4-dithiin, 1,4-oxathiin, oxazine, 1,3-oxazinone, morpholine, 1,3-dioxane, 1,3-dithiane, pyridazine, pyrazine, piperazine, 1,2,4-triazine, 1,3,5-triazine, 1,3,5-triaza-cyclohexane-2,4,6-trione;

**G** is a bridge between A and Z, **G'** is a bridge between NH and Z and **G''** is a bridge between E and Z, same or different, not necessary present, and are formed of one to ten moieties, each moiety being selected from the group consisting of phenylene, alkylene containing 1-12 carbon atoms, ethynediyl ( $-C\equiv C-$ ), ether ( $-O-$ ), thioether ( $-S-$ ), amide ( $-CO-NH-$  and  $-NH-CO-$  and  $-CO-NR'-$  and  $-NR'-CO-$ ), carbonyl ( $-CO-$ ), ester ( $-COO-$  and  $-OOC-$ ), disulfide ( $-S-S-$ ), diaza ( $-N=N-$ ), and tertiary amine ( $-N(R'')$ ), wherein  $R''$  represents an alkyl containing less than 5 carbon atoms;

$R'$  is  $-COOR'''$  where  $R'''$  is an alkyl of 1 to 4 carbon atoms, phenyl or benzyl, which phenyl or benzyl can be substituted or unsubstituted,

with a functional group which is bound to a solid phase,

b) removing the transient protecting group X and optionally other transient protecting groups attached to the solid phase,

c) reacting the labeling reactant of formula (I) with a deprotected functional group attached to the solid phase,

d) optionally repeating the steps b) to c) the number of times necessary to achieve the exact, predetermined number of labeling reactants of formula (I) bound to the solid phase,

e) cleaving the product obtained after the final step c) from the solid phase, converting  $R'$  to  $-COOH$ , removing the group or groups that have bound the product to the solid phase, and removing optional protecting groups, and

f) converting the product from step e) to a lanthanide chelate.

#### BRIEF DESCRIPTION OF THE DRAWINGS

##### [0013]

Figure 1 shows reversed phase HPLC trace of the oligopeptide conjugate R-E-S-Q-N-Y-P-I-V-X<sup>1</sup>-CONH<sub>2</sub> after deprotection (crude reaction mixture). X<sup>1</sup> is the deprotected labeling reagent 5 coupled to the COOH terminus of the oligopeptide, and

Figure 2 shows electron spray mass spectrum of the oligopeptide conjugate (major peak of the HPLC profile shown in Figure 1).

#### DETAILED DESCRIPTION OF THE INVENTION

[0014] The novel labeling reactants and novel labeling methods according to the present invention are particularly

suitable in the labeling of small molecules. Said small molecules are biospecific binding reactants including but not limited to antigens, haptens, peptides, receptor ligands, drugs or PNA oligomers which are used in specific bioaffinity based binding assays, such as immunoassays, DNA hybridization assays, receptor binding assays, immunocytochemical or immunohistochemical assays utilizing fluorometric or time-resolved fluorometric determination of the specific luminescence.

**[0015]** Particularly preferable transient protecting groups X are 2-(4-nitrophenylsulfonyl)ethoxycarbonyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, fluorenylmethyloxycarbonyl (Fmoc) or *tert*-butoxycarbonyl (Boc).

**[0016]** The substituent R" is preferably methyl, ethyl or *tert*-butyl.

**[0017]** The term "bivalent" in the definition of A shall mean a chemical group bound to two neighboring atoms.

**[0018]** The bivalent aromatic structure A is preferably selected from a group consisting of the structures disclosed in Scheme 1.

**[0019]** Most preferably, the labeling reactant is either 6-{N-[4-{2"-{2"-,6"-bis{N,N-bis[(*tert*-butoxycarbonyl)methyl]-aminomethyl)-4"-pyridyl]ethynyl}phenoxy)methylcarbonyl]amino}-2-{N-(fluorenylmethyloxycarbonyl)amino}hexanoic acid (**5**); 6-{N-[4-{2"-{1"-,3"-bis[(*tert*-butoxycarbonyl)methyl]amino-meth-5"-yl-2"-pyridyl}-1*H*-pyrazol-4"-yl)ethyl}phenoxy)methylcarbonyl]amino}-2-{N-(fluorenylmethyloxycarbonyl)-amino}hexanoic acid (**10**) or tetra(*tert*-butyl){2,2',2"-[4-[4-[N-(carboxymethyl)-N-[2-(4-methoxytritylamino)ethyl]aminocarbonylmethoxy]phenylethynyl]pyridine-2,6-di-yl]bis(methylenenitrilo)tetrakis(acetate)} (**13**).

**[0020]** According to a preferred embodiment of the invention the lanthanide chelate is a europium(III), terbium(III), samarium(III) or dysprosium(III) chelate.

**[0021]** The labeling of said biospecific binding reactants is performed by using solid-phase synthesis methods. It has appeared desirable to prepare suitable protected monomers that could be incorporated into said biospecific binding reactants during solid-phase synthesis. The labeling reactants of the present invention form a luminescent labeled biospecific binding reactant after releasing the product from solid support, deprotection and addition of a suitable lanthanide ion. The labeling technique of the present invention has many advantages over conventional labeling in liquid phase, such as high yields of easily purified products, straightforward routine automation leads to improved robustness, upscaling and repeatability of synthesis and labeling of biospecific binding reactant with an exact number of stable luminescent lanthanide chelates is possible. Although many of the advantages relate to solid-phase labeling, the compounds of the present invention can also be used in liquid phase labeling.

**[0022]** The labeling reactants and the methods of this invention combine several important features, such as:

- aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate,
- functional group or groups allowing efficient coupling of the compound to the said biospecific binding reactant without destroying its binding properties and decreasing the luminescence properties of the final labeled biospecific binding reactant.
- other functional groups are suitably protected against conditions and reactants used in solid-phase synthesis and it is easy to deprotect the functional groups during solid-phase synthesis and/or after the product has been released from the solid support.

**[0023]** It is, however, not absolutely necessary that the final biospecific binding reactant is bound to the solid phase already when the labeling reactant is attached to the functional group on the solid phase. The final biospecific binding reactant can be synthesized on the solid phase after the connection of the labeling reactant, or alternatively, the product comprising labeling reactants and optionally also spacer molecules, can be built up on the solid phase and then cleaved therefrom whereafter the cleaved product, before or after having been converted into a lanthanide chelate, is finally reacted in solution with the biospecific binding reactant.

**[0024]** According to one embodiment, a spacer molecule having a transient protecting group is reacted with a functional group on the solid phase and said transient protection group is removed prior to an introduction of the labeling reactant onto the solid phase. As an example of such a spacer molecule can be mentioned 6-aminohexanoic acid having the amino group protected with a transient protecting group, e.g. a group X as mentioned before.

**[0025]** The labeling reactant can be introduced into biomolecules with the aid of peptide synthesizer. The reagent is coupled to an amino tethered solid support or immobilized amino acid by e.g. carbodiimide chemistry described in Jones, J., The Chemical Synthesis of Peptides, Oxford University Press, Oxford, 1994, (i.e. the carboxylic acid function of the labeling reagent reacts with the amino group of the solid support or amino acid in the presence of an activator). When the condensation step is completed the transient amino protecting group of the labeling reagent is selectively removed while the material is still attached to the solid support (e.g. with piperidine in the case of Fmoc-protecting

group). Then second coupling of a labeling reagent or other reagent (amino acid, hapten) is performed as above. When the synthesis of the desired molecule is completed, the material is detached from the solid support and deprotected. Purification can be performed by HPLC techniques. Finally the purified ligand is converted to the corresponding lanthanide(III) chelate by addition of known amount of lanthanide(III) ion. (Mukkala, V.-M. et al. *Helv. Chim. Acta* 1993, 76, 1361 and M. Kwiatkowski, M. Samiotaki, U. Lamminmäki, V.-M. Mukkala and U. Landegren, *Nucleic Acid Res.*, 1994, Vol 22, 2604-2611).

[0026] The invention is further elucidated by the following examples. The structures and the synthetic routes employed in the experimental part are depicted in Schemes 2-5. Scheme 2 illustrates the synthetic procedures for the preparation of the labeling reagent 5. The experimental details are given in Examples 1-4. Scheme 3 illustrates synthetic procedures for the preparation of the labeling reagent 10. The experimental details are given in Examples 5-8. Scheme 4 illustrates the synthetic procedures for the preparation of the labeling reagent 13. The experimental details are given in Examples 9-11. Introduction of the labeling reagent 5 into an oligopeptide structure using machine assisted solid phase chemistry is described in Example 12. A solid phase method for the preparation of an estradiol conjugate tethered to four labeling reagents 5 is illustrated in Scheme 5. In order to increase the coupling efficiency, an aliphatic spacer molecule (6-aminohexanoic acid) is introduced between the labeling reagents. The experimental details are given in Example 13.

## EXPERIMENTAL PROCEDURES

[0027] Adsorption column chromatography was performed on columns packed with Kieselgel 60 (Merck). Analytical TLC was conducted on silica gel 60 F<sub>254</sub> plates (Merck) using the following solvent systems. System A: petroleum ether / ethyl acetate 5/2 (v/v); System B: CHCl<sub>3</sub>/MeOH 8/2 (v/v). NMR spectra were recorded on a Jeol LA-400 instrument operating at 399.8 and 105.0 MHz for <sup>1</sup>H and <sup>13</sup>C, respectively. TMS was used as an internal reference. Coupling constants are given in hertz. IR spectra were recorded on Perkin Elmer 2400 spectrophotometer.

### Example 1

The synthesis of methyl (4-ethynylphenoxy)acetate (2).

[0028] A solution of methyl (4-trimethylsilylethynylphenoxy)acetate (20.00 g, 76 mmol) in dichloromethane (400 mL) was deaerated with argon. Tetrabutylammonium fluoride 24.40 g, 91 mmol) was added, and the mixture was stirred for 30 min at ambient temperature. The solution was washed with 10% aqueous citric acid (200mL), water (5 x 400 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was dissolved in diethyl ether and passed through a column filled with silica gel. Evaporation of diethyl ether yielded the title compound as a solid (12.63 g, 83%) which was chromatographically and spectroscopically identical with material synthesized previously (Takalo, H., Hemmilä, I., Sutela, T. and Latva, M, 1996, *Helv. Chim. Acta*, 79, 789). R<sub>f</sub> (A) 0.50.

### Example 2

The synthesis of tetra(*tert*-butyl) 2,2',2'',2'''-[[4'-{(methoxycarbonylmethoxy)phenylethynyl} pyridine-2,6-diyl]bis (methylenenitrilo)]tetrakis(acetate) (3).

[0029] A mixture of tetra(*tert*-butyl) 2,2,2'',2'''-[(4-bromopyridine-2,6-diyl)bis-(methylenenitrilo)]tetrakis(acetate) (1) (2.32 g, 3.4 mmol), bis(triphenylphosphine)palladium(II) chloride (0.050 g, 70 mmol) and CuI (0.027 g, 140 mmol) in dry THF (10 mL) and dry triethylamine (10 mL) was deaerated with argon. Compound 2 (0.79 g, 4.1 mmol) was added and the mixture was stirred for 7 h at 50 °C. The cooled solution was filtered, the filtrate evaporated and redissolved in chloroform (150 mL). The solution was washed with water (3 x 150 mL), dried and evaporated. Purification on silica gel (System A) yielded the title compound as an oil (2.01 g, 75%). R<sub>f</sub> (A) 0.40. IR (film): 2212 (C≡C), 1750 (C=O), 1730(C=O). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.47 (36H, s), 3.50 (8H, s), 3.83 (3H, s), 4.02 (4H, s), 4.67 (2H, s), 6.90 (2H, d, J 9.0), 7.46 (2H, d, J 9.0), 7.60 (2H, s). <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 28.19, 52.36, 55.86, 59.73, 65.18, 81.25, 86.96, 92.89, 114.74, 115.90, 122.66, 132.68, 132.69, 133.44, 158.19, 168.98, 171.11.

### Example 3

The synthesis of 4-{2''',6'''-bis[N,N-bis[(*tert*-butoxycarbonyl)methyl]aminomethyl]-4'''-pyridyl}ethynyl} phenoxymethylacetic acid (4).

[0030] Compound 3 (1.98 g, 2.5 mmol) was dissolved in 0.25 M KOH (25 mL, 98 % EtOH, 2 % water) and the mixture



was stirred for 10 min at room temperature. The reaction was quenched by addition of 10% aqueous citric acid (100 mL). The resulting suspension was extracted with chloroform (100 mL). The organic layer was washed with citric acid (2 x 100 mL) and water (2 x 100 mL). Evaporation to dryness yielded the title compound as a solid (87%).  $R_f$  (B) 0.4. IR (film): 2209 (C≡C).  $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  1.41 (36H, s), 3.48 (8H, s), 3.89 (4H, s), 4.76 (2H, s), 6.99 (2H, d,  $J$  8.6), 7.52 (2H, d,  $J$  8.8), 7.52 (2H, s).

#### Example 4

The synthesis of 6-[N-[4-{2"-[2''',6'''-bis(*N,N*-bis[(*tert*-butoxycarbonyl)methyl] aminomethyl)-4'''-pyridyl]ethynyl] phenoxy methyl carbonyl] amino]-2[*N*-(fluorenylmethyloxycarbonyl) amino] hexanoic acid (**5**).

**[0031]** Compound **4** (0.75 g, 0.98 mmol) and *N*-hydroxysuccinimide (HO-Su) (0.13 g, 1.1 mmol) were dissolved in dry dioxane (30 mL). DCC (0.23 g, 1.1 mmol) predissolved in dry dioxane (5 mL) was added and the mixture was stirred for 4 h at room temperature and filtered on a solution of  $\alpha$ -Fmoc-lysine hydrochloride (0.88 g, 2.2 mmol) in the mixture of dry DMF (15 mL) and pyridine (5 mL). After 4 h, ethanol free chloroform (150 mL) was added and the solution was washed with 10% aq. citric acid (3 x 50 mL) and water (50 mL), dried (4 Å molecular sieves) and concentrated. Purification on silica gel (*System B*) yielded the title compound as a solid (56%). IR (film): 3315 (OH), 2210 (C≡C), 1681 (C=O).  $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  1.23 (2H, m), 1.40 (36H, s), 3.10 (2H, m), 3.30 (4H, m), 3.42 (8H, s), 3.89 (4H, s), 3.73 (1H, m), 4.24 (5H, m), 4.51 (2H, m), 7.01 (2H, d,  $J$  8.8), 7.32 (2H, t,  $J$  7.2), 7.40 (2H, t,  $J$  7.2), 7.51 (2H, s), 7.52 (2H, d,  $J$  8.8), 7.69 (2H, d,  $J$  7.5), 7.88 (2H, d,  $J$  7.5), 8.17 (1H, t,  $J$  6.3), 8.32 (1H, s).

#### Example 5

The synthesis of tetra(*tert*-butyl) 2,2',2'',2'''-{{6,6'-[4'-(methoxycarbonylmethoxy)phenylethynyl]-1*H*-pyrazol-1'',3''-diyl} bis(pyridine)-2,2'-diyl} bis-(methylenenitrilo)} tetrakis(acetate) (**7**).

**[0032]** A mixture of compound **6** (2.14 g, 2.5 mmol), bis(triphenylphosphine)palladium(II) chloride (35 mg, 50 mmol), and CuI (19 mg, 0.1 mmol) in dry DMF (12.5 mL) containing dry triethylamine (10 mL) was deaerated with argon. The mixture was stirred overnight at 35 °C. The solution was filtered and concentrated. The residue was dissolved in chloroform (100 mL), washed with water (3 x 100 mL), dried and concentrated. Purification was performed on silica gel column (*System A*). Yield was 1.12 g (45%).  $R_f$  (A) 0.2. IR (film): 1737 (C=O), 1143 (C-O, C=O).  $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  1.38 (18H, s), 1.42 (18H, s), 3.49 (4H, s), 3.52 (4H, s), 3.72 (3H, s), 4.00 (2H, s), 4.08 (2H, s), 4.86 (2H, s), 6.98 (2H, d,  $J$  8.9), 7.95 (1H, t,  $J$  7.8), 8.01 (1H, d,  $J$  7.1), 8.88 (1H, s).

#### Example 6

The synthesis of tetra(*tert*-butyl) 2,2',2'',2'''-{{6,6'-[2-[4'-(methoxycarbonylmethoxy)phenyl]ethyl]-1*H*-pyrazole-1'',3''-diyl} bis(pyridine)-2,2'-diyl} bis(methylenenitrilo)} tetrakis(acetate) (**8**).

**[0033]** Compound **7** (0.82 g, 0.88 mmol) was dissolved in dry methanol (60 mL). Pd/C (0.16 g, 10%) was added and the reaction was stirred for 4 h under hydrogen atmosphere. The solution was filtered through celite and evaporated. Yield was 0.57 g (70%).  $R_f$  (A) 0.3. IR (film): 1738 (C=O), 1143 (C-O).  $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  1.37 (18H, s), 1.41 (18H, s), 2.88 (2H, d,  $J$  8.4), 3.18 (2H, d,  $J$  8.4), 3.48 (4H, s), 3.49 (4H, s), 3.66 (3H, s), 3.98 (2H, s), 4.04 (2H, s), 4.73 (2H, s), 6.82 (2H, d,  $J$  8.4), 7.18 (2H, d,  $J$  8.4), 7.47 (1H, d,  $J$  7.6), 7.55 (1H, d,  $J$  7.6), 7.87 (1H, d,  $J$  7.8), 7.88 (1H, t,  $J$  7.6), 7.96 (1H, t,  $J$  7.8), 7.98 (1H, t,  $J$  7.6), 8.44 (1H, s).

#### Example 7

The synthesis of {4'-[2"-{1''',3'''-bis[(*tert*-butoxycarbonyl)methyl] aminometh-5'''-yl-2'''-pyridyl]-1*H*-pyrazol-4'''-yl} ethylphenoxy} acetic acid (**9**).

**[0034]** The title compound (**9**) was synthesized as described above for **4** in Example 2 by using compound **8** (0.57 g, 0.62 mmol) as the starting material. Purification was performed on silica gel (petroleum ether / ethyl acetate 5/2 → MeOH/CHCl<sub>3</sub> 2/8). Yield was 100%.  $R_f$  (B) 0.95.  $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  1.37 (18H, s), 1.41 (18H, s), 2.84 (2H, d,  $J$  8.4), 3.15 (2H, d,  $J$  8.4), 3.47 (4H, s), 3.48 (4H, s), 3.98 (2H, s), 4.03 (2H, s), 4.17 (2H, s), 6.74 (2H, d,  $J$  8.4), 7.11 (2H, d,  $J$  8.4), 7.46 (1H, d,  $J$  7.8), 7.55 (1H, d,  $J$  6.9), 7.87 (1H, d,  $J$  7.8), 7.88 (1H, t,  $J$  7.9), 7.95 (1H, d,  $J$  6.9), 7.98 (1H, t,  $J$  7.8), 8.44 (1H, s).

Example 8

The synthesis of 6-{N-[4'-(2''-{1'''-3'''-bis[(*tert*-butoxycarbonyl)methyl]aminometh-5''''-yl-2''''-pyridyl)-1*H*-pyrazol-4''''-yl)ethyl]phenoxyethylcarbonyl}-amino-2-[*N*-(fluorenylmethyloxycarbonyl) amino]hexanoic acid (**10**).

**[0035]** The title compound (**10**) was synthesized as described in Example 4 for **5** by using **9** as the starting material. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 1.05 (1H, m), 1.20 (2H, m), 1.36 (18H, s), 1.40 (18H, s), 1.60 (2H, s), 1.71 (2H, m), 2.86 (2H, t, *J* 7.9), 3.10 (2H, m), 3.18 (2H, m), 3.30 (4H, s), 3.38 (4H, s), 3.80 (1H, m), 3.97 (2H, s), 4.03 (2H, s), 4.23 (4H, m), 4.39 (2H, s), 6.85 (2H, d, *J* 8.6), 7.17 (2H, d, *J* 8.6), 7.31 (2H, t, *J* 7.4), 7.40 (2H, t, *J* 7.4), 7.47 (1H, d, *J* 7.7), 7.55 (1H, d, *J* 7.7), 7.70 (2H, d, *J* 7.3), 7.88 (1H, d, *J* 7.6), 7.99 (1H, d, *J* 7.8), 8.44 (1H, s).

Example 9

The synthesis of methyl *N*-[2-{4-methoxytrityl}amino]ethylglycinate (**11**).

**[0036]** *N*1-[4-methoxytrityl]-1,2-diaminoethane (4.69 g, 14.2 mmol) was dissolved in dry THF (20 mL). Dry TEA (2 mL) and methyl bromoacetate (1.58 mL, 14.2 mmol) were added, and the mixture was stirred for 1 h at room temperature, after which the precipitation formed was filtered. After concentration the filtrate was dissolved in ethyl acetate, washed with sat. Na<sub>2</sub>SO<sub>4</sub>, water and dried. Purification on silica gel (eluent PE:EA:TEA 5:1:1) yielded the title compound as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.93 (1H, br), 2.26 (2H, t, *J* 5.6), 2.72 (2H, t, *J* 5.6), 3.34 (2H, s), 3.69 (3H, s), 3.74 (3H, s), 6.80 (2H, d, *J* 8.6), 7.14 (2H, t), 7.24 (4H, t), 7.36 (2H, d), 7.46 (4H, t).

Example 10

The synthesis of tetra(*tert*-butyl) {2,2',2'',2'''-[4-[4-[*N*-(methoxycarbonylmethyl)-*N*-[2-(4-methoxytritylamino)ethyl]aminocarbonylmethoxy]phenylethynyl]pyridine-2,6-diyl]bis(methylenenitrilo)tetrakis(acetate)} (**12**)

**[0037]** Compound **4** (0.5 g, 0.65 mmol), compound **11** (0.26 g, 0.65 mmol) and *N*-hydroxysuccinimide (115 mg, 1.0 mmol) were dissolved in dry acetonitrile. DCC (134 mg, 0.65 mmol; predissolved in dry acetonitrile) was added and the mixture was stirred overnight at room temperature. DCU formed was filtered and the filtrate was concentrated in vacuo. Purification on silica gel yielded **12**. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.41 (36H, s), 2.27 (2H, m), 2.61 (2H, m), 3.38 (2H, s), 3.47 (8H, s), 3.68 (3H, s), 3.76 (3H, s), 4.00 (4H, s), 6.77 (2H, d, *J* 8.8), 6.89 (2H, d, *J* 8.8), 7.41-7.14 (14H, m), 7.51 (2H, s).

Example 11

The synthesis of tetra(*tert*-butyl) {2,2',2'',2'''-[4-[4-[*N*-(carboxymethyl)-*N*-[2-(4-methoxytritylamino)ethyl]aminocarbonylmethoxy]phenylethynyl]pyridine-2,6-diyl]bis(methylenenitrilo)tetrakis(acetate)} (**13**).

**[0038]** Compound **12** was hydrolyzed and purified as described in Example 3. Yield was 65%. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.39 (36H, s), 2.26 (2H, m), 2.61 (2H, m), 3.38 (2H, s), 3.47 (8H, s), 3.70 (3H, s), 3.78 (3H, s), 4.00 (4H, s), 6.77 (2H, d, *J* 8.8), 6.89 (2H, d, *J* 8.8), 7.42-7.17 (14H, m), 7.54 (2H, s).

Example 12

Introduction of the labeling reagent into oligopeptides.

**[0039]** In order to demonstrate the applicability of the ligands prepared to oligopeptide labeling, a sequence R-E-S-Q-N-Y-P-I-V-X<sup>1</sup>-CONH<sub>2</sub> was synthesized on a Perkin Elmer 433A peptide synthesizer (5 mmol scale) using standard Fmoc chemistry. The ligand **5** was introduced in the first coupling step using a prolonged coupling time (1 h instead of 30 min). When the chain assembly was completed, the oligopeptide was deprotected and purified by HPLC. A typical HPLC trace (a crude reaction mixture) is shown in Figure 1. The product was confirmed by electron-spray mass spectrometry (Figure 2). Finally the purified material was converted into the corresponding europium(III) chelate using the standard protocol.

Example 13

Labeling of estradiol with four europium(III) chelates.

5   **[0040]** The labeling reagent 5 was introduced into an amino derivatized solid support as described in Example 12. After removal of the  $\alpha$ -Fmoc protecting group 6-Fmoc-aminohexanoic acid was coupled. These cycles were repeated three times and an estradiol derivative tethered to a carboxylic acid linker was introduced as the last coupling step. Deprotection and purification were performed as described in Example 12.

10   **[0041]** It will be appreciated that the methods of the present invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will be apparent for the specialist in the field that other embodiments exist and do not depart from the spirit of the invention. Thus, the described embodiments are illustrative and should not be construed as restrictive.

15

20

25

30

35

40

45


50

55

## Scheme 1


5

10



15


20



25



30




35

40




45

50




55


CC(C)(C)C(=O)OCCN(CCN(CCC(C)(C)C)CC(C)(C)C)c1ccc(Br)cc1 (1)
   
 $\xrightarrow{\text{PdCl}_2(\text{Ph}_3\text{P})_2/\text{CuI}}$ 
CC(C)(C)C(=O)OCCN(CCN(CCC(C)(C)C)CC(C)(C)C)c1ccc(C#Cc2ccc(OCC(=O)OC)cc2)cc1 (2)
   
 $\xrightarrow{\text{1. HO-Su/DCC}}$ 
 $\xrightarrow{\text{2. } \alpha\text{-Fmoc-L-Lys}}$ 
CC(C)(C)C(=O)OCCN(CCN(CCC(C)(C)C)CC(C)(C)C)c1ccc(C#Cc2ccc(OCC(=O)OCC(=O)OC)cc2)cc1 (3)
   
 $\xrightarrow{\text{0.25 M KOH}}$ 
 $\xrightarrow{\text{98\% MeOH}}$ 
 $\xrightarrow{\text{2\% H}_2\text{O}}$ 
CC(C)(C)C(=O)OCCN(CCN(CCC(C)(C)C)CC(C)(C)C)c1ccc(C#Cc2ccc(OCC(=O)O)cc2)cc1 (4)



## Scheme 3



## Scheme 4




**Scheme 3**

The scheme illustrates the synthesis of a complex molecule, likely a peptide or protein derivative, through iterative cycles of coupling and deblocking. The starting material is a steroid derivative with a hydroxyl group and an amine group. The reaction proceeds through 4 cycles of coupling using Fmoc-aminohexanoic acid, followed by 1 cycle of deblocking using  $\text{EuCl}_3$ . The final product is a complex molecule featuring a steroid core, a peptide backbone, and a haptens (protected hapten) attached via a linker. The structure shows the coupling of the steroid derivative with the haptens, resulting in a complex with multiple amine and carboxylate groups, and a central metal ion ( $\text{Eu}^{3+}$ ) coordinated to the complex.

1. A labeling reactant of formula (I) suitable for labeling of a biospecific binding reactant using solid-phase synthesis





wherein


15 **A** is a bivalent aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate;

20 **R** is -Z(G'-NH-X)G"-E, where

**X** is a transient protecting group, where said group is e.g. 2-(4-nitrophenylsulfonyl)ethoxycarbonyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, *tert*-butoxycarbonyl, fluorenylmethyloxycarbonyl;

25 **E** is a carboxylic acid, its salt, active ester or halide, where the said ester is e.g. an *N*-hydroxysuccinimido, *o*-, *p*- or *m*-nitrophenol, 2,4-dinitrophenol or pentafluorophenol, and the said halide is fluoride, bromide or chloride;

**Z** is the bridge point and is formed from



40 or trivalent derivatives, substituted or unsubstituted, of cyclohexane, cyclohexene, cyclohexadiene, benzene, cyclopentane, cyclopentene, cyclopentadiene, cyclobutane, cyclobutene, cyclobutadiene, aziridine, diaziridine, oxetane, thietane, azete, azetidine, 1,2-dihydro-1,2-diazete, 1,2-diazetidine, furan, tetrahydrofuran, thiophene, 2,5-dihydrothiophene, thiolane, selenophene, pyrrole, pyrrolidine, phosphole, 1,3-dioxolane, 1,2-dithiole, 1,2-thiolane, 1,3-dithiole, 1,3-dithiolane, oxazole, 4,5-dihydrooxazole, isoxazole, 4,5-dihydroisoxazole, 2,3-dihydroisoxazole, thiazole, isothiazole, imidazole, imidazolidine, pyrazole, 4,5-dihydropyrazole, pyrazolidine, triazole, pyran, pyran-2-one, 3,4-dihydro-2H-pyran, tetrahydropyran, 4H-pyran, pyran-4-one, pyridine, pyridone, piperidine, phosphabenzene, 1,4-dioxin, 1,4-dithiin, 1,4-oxathiin, oxazine, 1,3-oxazinone, morpholine, 1,3-dioxane, 1,3-dithiane, pyridazine, pyrimidine, pyrazine, piperazine, 1,2,4-triazine, 1,3,5-triazine, 1,3,5-triaza-cyclohexane-2,4,6-trione;


50 **G** is a bridge between A and Z, **G'** is a bridge between NH and Z and **G"** is a bridge between E and Z, same or different, not necessary present, and are formed of one to ten moieties, each moiety being selected from the group consisting of phenylene, alkylene containing 1-12 carbon atoms, ethynediyl (-C≡C-), ether (-O-), thioether (-S-), amide (-CO-NH- and -NH-CO- and -CO-NR'- and -NR'-CO-), carbonyl (-CO-), ester (-COO- and -OOC-), disulfide (-S-S-), diaza (-N=N-), and tertiary amine (-N(R'')-), wherein R'' represents an alkyl containing less than 5 carbon atoms;

55 **R'** is -COOR''' where R''' is an alkyl of 1 to 4 carbon atoms, phenyl or benzyl, which phenyl or benzyl can be substituted or unsubstituted.

2. The labeling reactant according to claim 1 wherein X is fluorenylmethyloxycarbonyl (Fmoc), *tert*-butoxycarbonyl

(Boc) or 4-methoxytrityl.

3. The labeling reactant according to claim 1 wherein R<sup>'''</sup> is methyl, ethyl or *tert*-butyl.
- 5 4. The labeling reactant according to claim 1 wherein the biospecific binding reactant is selected from a group consisting of an antigen, a hapten, a peptide, a receptor ligand, a drug or PNA oligomer.
5. The labeling reactant according to claim 1 wherein -A- is a bivalent aromatic structure selected from a group consisting of the structures disclosed in Scheme 1.
- 10 6. The labeling reactant according to claim 1 selected from the group consisting of 6-{*N*-(4-(2''-(2''',6'''-bis(*N,N*-bis[(*tert*-butoxycarbonyl)methyl]aminomethyl)-4'''-pyridyl)ethynyl)phenoxy)methylcarbonyl]amino)-2-[*N*-(fluorenylmethyloxycarbonyl)amino]hexanoic acid (**5**), 6-{*N*-(4'-(2''-(1''',3'''-bis[(*tert*-butoxycarbonyl)-methyl]aminomethyl)-5'''-yl)-2'''-pyridyl)-1*H*-pyrazol-4'''-yl)ethyl}-phenoxy)methylcarbonyl]amino)-2-[*N*-(fluorenylmethyloxycarbonyl)amino]hexanoic acid (**10**) and tetra(*tert*-butyl){2,2',2'',2'''-[[4-[4-[*N*-(carboxymethyl)-*N*-(2-(4-methoxytritylamino)ethyl)aminocarbonylmethoxy]phenylethynyl]pyridine-2,6-diyl]bis(methylenenitrilo)tetrakis(acetate)]} (**13**).
- 15 7. A method of labeling a biospecific binding reactant in a solid-phase synthesis, said method comprising the steps of
  - 20 a) reacting a labeling reactant of formula (I)



wherein


**A** is a bivalent aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate;

**R** is -Z(G'-NH-X)G"-E, where

**X** is a transient protecting group, where said group is e.g. 2-(4-nitrophenylsulfonyl)ethoxycarbonyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, *tert*-butoxycarbonyl, fluorenylmethyloxycarbonyl;

**E** is a carboxylic acid, its salt, active ester or halide, where the said ester is e.g. an *N*-hydroxysuccinimido, *o*-, *p*- or *m*-nitrophenol, 2,4-dinitrophenol or pentafluorophenol, and the said halide is fluoride, bromide or chloride;

**Z** is the bridge point and is formed from



or trivalent derivatives, substituted or unsubstituted, of cyclohexane, cyclohexene, cyclohexadiene, benzene, cyclopentane, cyclopentene, cyclopentadiene, cyclobutane, cyclobutene, cyclobutadiene, aziridine, diaziridine, oxetane, thietane, azete, azetidine, 1,2-dihydro-1,2-diazete, 1,2-diazetidene, furan, tetrahydrofuran, thiophene, 2,5-dihydrothiophene, thiolane, selenophene, pyrrole, pyrrolidine, phosphole, 1,3-dioxolane, 1,2-dithiole, 1,2-thiolane, 1,3-dithiole, 1,3-dithiolane, oxazole, 4,5-dihydrooxazole, isoxazole, 4,5-dihydroisoxazole, 2,3-dihydroisoxazole, thiazole, isothiazole, imidazole, imidazolidine, pyrazole, 4,5-dihydropyrazole, pyrazolidine, triazole, pyran, pyran-2-one, 3,4-dihydro-2H-pyran, tetrahydropyran, 4H-pyran, pyran-4-one, pyridine, pyridone, piperidine, phosphabenzene, 1,4-dioxin, 1,4-dithiin, 1,4-oxathiin, oxazine, 1,3-oxazinone, morpholine, 1,3-dioxane, 1,3-dithiane, pyridazine, pyrimidine, pyrazine, piperazine, 1,2,4-triazine, 1,3,5-triazine, 1,3,5-triaza-cyclohexane-2,4,6-trione;

**G** is a bridge between A and Z, **G'** is a bridge between NH and Z and **G''** is a bridge between E and Z, same or different, not necessary present, and are formed of one to ten moieties, each moiety being selected from the group consisting of phenylene, alkylene containing 1-12 carbon atoms, ethynediyl ( $-C\equiv C-$ ), ether ( $-O-$ ), thioether ( $-S-$ ), amide ( $-CO-NH-$  and  $-NH-CO-$  and  $-CO-NR'-$  and  $-NR'-CO-$ ), carbonyl ( $-CO-$ ), ester ( $-COO-$  and  $-OOC-$ ), disulfide ( $-S-S-$ ), diaza ( $-N=N-$ ), and tertiary amine ( $-N(R'')$ ), wherein  $R''$  represents an alkyl containing less than 5 carbon atoms:

**R'** is  $-COOR'''$  where  $R'''$  is an alkyl of 1 to 4 carbon atoms, phenyl or benzyl, which phenyl or benzyl can be substituted or unsubstituted,

with a functional group which is bound to a solid phase,

b) cleaving the product obtained in step a) from the solid phase, converting  $R'$  to  $-COOH$ , and removing optional protection groups,

c) converting the product from step b) to a lanthanide chelate, and

d) in case the functional group mentioned in step a) is not part of the final biospecific binding specific reactant, attaching said biospecific binding reactant to the lanthanide chelate obtained in step c).

8. The method according to claim 7 wherein X in the labeling reactant is fluorenylmethyloxycarbonyl (Fmoc), *tert*-butoxycarbonyl (Boc) or 4-methoxytrityl.

9. The method according to claim 7 wherein  $R'''$  in the labeling reactant is methyl, ethyl or *tert*-butyl.


10. The method according to claim 7 wherein the biospecific binding reactant is selected from a group consisting of an antigen, a hapten, a peptide, a receptor ligand, a drug or PNA oligomer.

11. The method according to claim 7 wherein -A- in the labeling reactant is a bivalent aromatic structure selected from a group consisting of the structures disclosed in Scheme 1.

12. The method according to claim 7 wherein the labeling reactant is selected from the group consisting of 6-{N-[4-{2''-[2''',6'''-bis({N,N-bis({*tert*-butoxycarbonyl)methyl} aminomethyl)-4'''-pyridyl]ethynyl}phenoxy)methyl-carbonyl)-amino}-2-[N-(fluorenylmethyloxycarbonyl)amino]hexanoic acid (**5**), 6-{N-[4'-{2''-[1''',3'''-bis({*tert*-butoxycarbonyl)-methyl]aminomethyl-5'''-yl-2'''-pyridyl]-1H-pyrazol-4'''-yl]ethyl}phenoxy)methyl-carbonyl)amino-2-[N-(fluorenylmethyloxycarbonyl)amino]hexanoic acid (**10**) and tetra(*tert*-butyl) {2,2',2'',2'''-[4-[4-N-(carboxymethyl)-N-[2-(4-methoxytritylamino)ethyl]aminocarbonylmethoxy]phenylethynyl]pyridine-2,6-diyl]bis(methylenenitrilo)tetraakis(acetate)} (**13**).

13. A method of labeling a biospecific binding reactant with an exact, predetermined number of labeling reactants of formula (I) in a solid-phase synthesis, said method comprising the steps of

a) reacting a labeling reactant of formula (I)



15 wherein


**A** is a bivalent aromatic structure capable of absorbing light or energy and transferring the excitation energy to a lanthanide ion after the product made by the said solid-phase synthesis has been released from the used solid support, deprotected and converted to a lanthanide chelate;

20 **R** is - Z(G'-NH-X)G"- E, where

**X** is a transient protecting group, where said group is e.g. 2-(4-nitrophenylsulfonyl)ethoxycarbonyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, tert-butoxycarbonyl, fluorenylmethyloxycarbonyl;

25 **E** is a carboxylic acid, its salt, active ester or halide, where the said ester is e.g. an N-hydroxysuccinimido, *o*-, *p*- or *m*-nitrophenol, 2,4-dinitrophenol or pentafluorophenol, and the said halide is fluoride, bromide or chloride;

30 **Z** is the bridge point and is formed from



40 or trivalent derivatives, substituted or unsubstituted, of cyclohexane, cyclohexene, cyclohexadiene, benzene, cyclopentane, cyclopentene, cyclopentadiene, cyclobutane, cyclobutene, cyclobutadiene, aziridine, diaziridine, oxetane, thietane, azete, azetidine, 1,2-dihydro-1,2-diazete, 1,2-diazetidine, furan, tetrahydrofuran, thiophene, 2,5-dihydrothiophene, thiolane, selenophene, pyrrole, pyrrolidine, phosphole, 1,3-dioxolane, 1,2-dithiole, 1,2-thiolane, 1,3-dithiole, 1,3-dithiolane, oxazole, 4,5-dihydrooxazole, isoxazole, 4,5-dihydroisoxazole, 2,3-dihydroisoxazole, thiazole, isothiazole, imidazole, imidazolidine, pyrazole, 4,5-dihydropyrazole, pyrazolidine, triazole, pyran, pyran-2-one, 3,4-dihydro-2H-pyran, tetrahydropyran, 4H-pyran, pyran-4-one, pyridine, pyridone, piperidine, phosphabenzene, 1,4-dioxin, 1,4-dithiin, 1,4-oxathiin, oxazine, 1,3-oxazinone, morpholine, 1,3-dioxane, 1,3-dithiane, pyridazine, pyrimidine, pyrazine, piperazine, 1,2,4-triazine, 1,3,5-triazine, 1,3,5-triaza-cyclohexane-2,4,6-trione;

50 **G** is a bridge between A and Z, **G'** is a bridge between NH and Z and **G"** is a bridge between E and Z, same or different, not necessary present, and are formed of one to ten moieties, each moiety being selected from the group consisting of phenylene, alkylene containing 1-12 carbon atoms, ethynediyl (-C≡C-), ether (-O-), thioether (-S-), amide (-CO-NH- and -NH-CO- and -CO-NR'- and -NR'-CO-), carbonyl (-CO-), ester (-COO- and -OOC-), disulfide (-S-S-), diaza (-N=N-), and tertiary amine (-N(R")-), wherein R" represents an alkyl containing less than 5 carbon atoms;

55 **R'** is -COOR''' where R''' is an alkyl of 1 to 4 carbon atoms, phenyl or benzyl, which phenyl or benzyl can be substituted or unsubstituted,

with a functional group which is bound to a solid phase.

b) removing the transient protecting group X and optionally other transient protecting groups attached to the solid phase.

c) reacting the labeling reactant of formula (I) to a deprotected functional group attached to the solid phase,

d) optionally repeating the steps b) to c) the number of times necessary to achieve the exact, predetermined number of labeling reactants of formula (I) bound to the solid phase,

e) cleaving the product obtained after the final step c) from the solid phase, converting R' to -COOH, removing the group or groups that have bound the product to the solid phase, and removing optional protecting groups, and

f) converting the product from step e) to a lanthanide chelate.

14. The method according to claim 13 wherein step f) is followed by reacting the biospecific binding reactant with the lanthanide chelate obtained in step f), to obtain the labeled biospecific binding reactant.

15. The method according to claim 13 wherein the biospecific binding reactant is reacted with the product containing the labeling reactants while said product still is bound to the solid phase.

16. The method according to claim 13 wherein the biospecific binding reactant is built up in the solid phase synthesis.

17. The method according to claim 13 wherein a spacer molecule having a transient protecting group is reacted with a functional group on the solid phase and said transient protection group is removed prior to an introduction of the labeling reactant onto the solid phase.


18. The method according to claim 13 wherein X in the labeling reactant is fluorenylmethyloxycarbonyl (Fmoc), *tert*-butoxycarbonyl (Boc) or 4-methoxytrityl.

19. The method according to claim 13 wherein R''' in the labeling reactant is methyl, ethyl or *tert*-butyl.

20. The method according to claim 13 wherein the biospecific binding reactant is selected from a group consisting of an antigen, a hapten, a peptide, a receptor ligand, a drug or PNA oligomer.

21. The method according to claim 13 wherein A in the labeling reactant is a bivalent aromatic structure selected from a group consisting of the structures disclosed in Scheme 1.

22. The method according to claim 13 wherein the labeling reactant is selected from the group consisting of 6-{N-[4-{2''-{2'''-6'''-bis{N,N-bis[(*tert*-butoxycarbonyl)methyl]-aminomethyl]-4'''-pyridyl}ethynyl}phenoxy)methylcarbonyl}amino]-2-{N-(fluorenylmethyloxycarbonyl)amino}hexanoic acid (**5**), 6-{N-[4'-{2''-{1'''-3'''-bis[(*tert*-butoxycarbonyl)-methyl]aminometh-5'''-yl-2'''-pyridyl}-1 H-pyrazol-4'''-yl}ethyl}phenoxy)methylcarbonyl}amino-2-[N-(fluorenylmethyloxycarbonyl)-amino]hexanoic acid (**10**) and tetra(*tert*-butyl){2,2',2''-[[4-[4-N-(carboxymethyl)-N-[2-(4-methoxytritylamino)ethyl]aminocarbonylmethoxy]phenylethynyl]pyridine-2,6-diyl]bis(methylenenitrilo)tetrakis (acetate)} (**13**).



**FIG. 1**




FIG. 2



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 99 66 0100

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D,A	EP 0 770 610 A (WALLAC OY) 2 May 1997 (1997-05-02) * claims *	1	C07D213/38 C07D401/14 G01N33/48
A	WO 93 11433 A (WALLAC OY) 10 June 1993 (1993-06-10) * claims *	1	
D,A	DE 44 30 998 A (BOEHRINGER MANNHEIM GMBH) 1 February 1996 (1996-02-01) * claims *	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07D G01N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 21 September 1999	Examiner Van Bijlen, H
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, not published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPC FORM 1503 03 82 (P4/C01)



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 66 0100

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

21-09-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 770610 A	02-05-1997	US 5859215 A	12-01-1999
WO 9311433 A	10-06-1993	DE 69131206 D	10-06-1999
		EP 0626068 A	30-11-1994
		US 5571897 A	05-11-1996
DE 4430998 A	01-02-1996	AU 682278 B	25-09-1997
		AU 3164995 A	22-02-1996
		AU 689626 B	02-04-1998
		AU 3165095 A	22-02-1996
		AU 690315 B	23-04-1998
		AU 3220595 A	22-02-1996
		CA 2195648 A	08-02-1996
		CA 2195752 A	08-02-1996
		CN 1134154 A	23-10-1996
		CN 1152923 A	25-06-1997
		WO 9603650 A	08-02-1996
		WO 9603651 A	08-02-1996
		WO 9603409 A	08-02-1996
		WO 9603410 A	08-02-1996
		EP 0772616 A	14-05-1997
		EP 0774119 A	21-05-1997
		EP 0774120 A	21-05-1997
		EP 0720614 A	10-07-1996
		FI 961349 A	25-03-1996
		FI 970299 A	24-01-1997
		FI 970300 A	24-01-1997
		JP 9508473 T	26-08-1997
		JP 10506708 T	30-06-1998
		JP 2771900 B	02-07-1998
		JP 8509995 T	22-10-1996
		JP 10503485 T	31-03-1998
		NO 961161 A	21-03-1996
		NO 970130 A	10-01-1997
		NO 970197 A	16-01-1997
		NZ 290810 A	26-01-1998
		NZ 290811 A	19-12-1997
		NZ 291154 A	24-11-1997
		AU 688953 B	19-03-1998
		AU 3220495 A	22-02-1996
		AU 684992 B	08-01-1998
		AU 3220695 A	22-02-1996
		CA 2195753 A	08-02-1996
		CN 1130910 A	11-09-1996
		CN 1157655 A	20-08-1997
		DE 4430972 A	01-02-1996

EPC FORM P0459

For more details about this annex, see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 66 0100

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

21-09-1999

Patent document cited in search report	Publication date	Patent family members)	Publication date
DE 4430998 A		DE 4430973 A	01-02-1996
		DE 4439345 A	01-02-1996
		DE 4439346 A	01-02-1996
		DE 4439347 A	01-02-1996
		WO 9603652 A	08-02-1996
		WO 9603423 A	08-02-1996
		EP 0772774 A	14-05-1997
		EP 0723551 A	31-07-1996
		FI 961350 A	25-03-1996
<hr/>			

EPO FORM P/459

For more details about this annex see Official Journal of the European Patent Office, No. 12/82